GEOHELMINTHS EGGS CONTAMINATION OF SANDPITS IN VLADIVOSTOK, RUSSIA.


Moskvina Tatyana Vladimirovna*

Bartkova Albina Dmitrievna¹


Ermolenko Alexey Vasil'evich²

The most widespread geohelminths

? Toxocara canis life cycle



? Ascaris lumbricoides life cycle



Materials and methods

? Totally 60 samples from five district were collected and analyzed. Sand samples were analyzed using Romanenko's method






NaNO₃

Results



Egg with larva

Toxocara spp.eggs



Ascaris sp.egg



Toxascaris leonina egg



Results

Table 1 Parasites found in soil samples

Parasites found	Number of soil samples	Min-max number of eggs/25
		gr
Toxascaris leonina	2	1-23
Toxocara	14	1-6
Ascaris sp.	7	1-3

Table 2 Results of sandpits samples analyses in each district

District	Total number of	Number of	Min-max	Sovetskij
	collected samples	positive samples	number of	
			eggs/25 g	Pervorechenskij
Pervorechenskij	12	3	3-5	
Sovetskij	11	3	1-3	Frunzenskij
Pervomajskij	16	5	1-26	Pervomajskij
Frunzenskij	11	4	1-3	
Leninskii'	10	3	2-18	
		<u> </u>	1	

Control and prevention





Thanks' for your attention