

Как подготовить стендовый доклад?

Что такое стендовый доклад?

- это результаты научного исследования, оформленные в виде большого бумажного плаката (постера).
- **Постер-доклад** – достаточно новая для России, но хорошо себя зарекомендовавшая форма представления результатов научного исследования на крупной конференции. Автор демонстрирует свой материал в письменной форме, коллеги имеют возможность ознакомиться с ним и задать вопросы непосредственно в процессе знакомства с данными.

Примеры:

RNA interference in mammalian cells using low siRNA concentrations



Jörg Drenth*, Silvia Magyar*, Anja Grewe*, Cornelia Schmidt*, Peter Hahn*, Dong Liang*, Subu Yerramilil*, Eric Lader*, Wolfgang Bieleke*, and Jie Kang*

* QIAGEN GmbH, Hilden, Germany; * QIAGEN Sciences, Germantown, MD, USA

Introduction

The use of short interfering RNA (siRNA) for knockdown of gene expression has become a powerful tool in molecular and cell biology. Some applications require the use of low siRNA concentrations (less than 5 nM), for example, to determine the possibility of nonspecific effects.

We have developed a transfection reagent, HiPerfect Transfection Reagent, which allows efficient gene knockdown with siRNA concentrations from 1 nM to 10 nM, depending on the cell type and siRNA used. HiPerfect Transfection Reagent has been tested and validated for many cell types, including primary cells. Efficient knockdown in primary cells demonstrates that HiPerfect Transfection Reagent works for significantly lower siRNA concentrations.



Highly effective knockdown of CDC2 expression with low siRNA concentrations

Comparison of Knockdown Efficiency Using HiPerfect Transfection Reagent and Reagent L





Figure 1: CDC2 expression was measured in a range of concentrations of siRNA. HiPerfect Transfection Reagent showed significantly higher knockdown efficiency than Reagent L. The knockdown efficiency was measured by relative expression of CDC2 mRNA. The knockdown efficiency was measured by relative expression of CDC2 mRNA. The knockdown efficiency was measured by relative expression of CDC2 mRNA.



HiPerfect Transfection Reagent from QIAGEN allowed highly efficient CDC2 knockdown with siRNA concentrations as low as 1 nM. In contrast, Reagent L from another supplier provided less efficient knockdown of all concentrations tested. For concentrations lower than 5 nM, knockdown of only 15% or less was observed.

Transfection and knockdown in a wide range of cell types

A wide range of cell types have been successfully transfected using HiPerfect Transfection Reagent. For an up-to-date list of cell types and more detailed information go to www.qiagen.com/transfectioninfo.

A Wide Range of Successfully Transfected Cells



Figure 1: The transfection of cells in a wide range of cell types using HiPerfect Transfection Reagent.

Rapid, efficient lamin A/C knockdown in human primary cells

Lamin A/C Knockdown Using the Fast-Forward Protocol with Low siRNA Concentration




Figure 2: Lamin A/C knockdown was measured in a range of concentrations of siRNA. The Fast-Forward Protocol showed significantly higher knockdown efficiency than the standard protocol. The knockdown efficiency was measured by relative expression of lamin A/C mRNA. The knockdown efficiency was measured by relative expression of lamin A/C mRNA.




Figure 3: Phase contrast picture of transfected cells. The image shows cells with varying degrees of transfection, with the most significant transfection observed in cells treated with the Fast-Forward Protocol.

Summary

- HiPerfect Transfection Reagent allows gene silencing using as little as 1 nM siRNA.
- Transfection of low siRNA concentrations may be necessary to avoid off-target effects. Using HiPerfect Transfection Reagent means that effective knockdown can be achieved with very low siRNA concentrations.
- HiPerfect Transfection Reagent has been tested for a range of cell types.
- Many cell types and primary cells have been successfully transfected using low siRNA concentrations and HiPerfect Transfection Reagent. For an up-to-date list, visit www.qiagen.com/transfectioninfo.
- The Fast-Forward Protocol has been developed for rapid transfection.
- The Fast-Forward Protocol, cell seeding and siRNA transfection are carried out on the same day, saving time and effort. The Fast-Forward Protocol is available at www.qiagen.com/fastforward.
- The reverse transfection protocol is ideal for RNAi screening.
- The reverse transfection protocol can be easily extended which is particularly useful for high-throughput applications. The reverse transfection protocol is available at www.qiagen.com/fastforward.

Transfection Reagent, HiPerfect Transfection Reagent, Fast-Forward Protocol, and HiPerfect Transfection Reagent are trademarks of QIAGEN. All other trademarks are the property of their respective owners. © 2008 QIAGEN. All rights reserved.

HiPerfect Transfection Reagent Allows Effective Uptake of Low Amounts of Alexa Fluor® 488 Labeled siRNA

Reversibly labeled siRNA Shows Highly Efficient Uptake into Cells





Figure 1: Reversible labeling of siRNA with Alexa Fluor 488 allows efficient uptake into cells. The image shows cells with varying degrees of uptake, with the most significant uptake observed in cells treated with HiPerfect Transfection Reagent.

WWW.QIAGEN.COM

Biomarkers and Metabolomics: Practical Implication.

Vladimir V. Tolstikov

UC Davis Genome Center, Davis, CA




Department of Internal Medicine
UC DAVIS HEALTH SYSTEM

Robert H. Weiss, M.D.
Division of Nephrology & Cancer Center

Phase I Aim: Find potential small molecule biomarkers for metastatic kidney cancer (RCC). Metabolomics pilot study. Proof of the concept. Phase I is completed.

Phase II Aim: Use Phase I proof of concept methodology to carry out next set of studies: Run Metabolomics study on large group cancer patients and volunteers. Identify the most prominent biomarkers suitable for RCC diagnostic test development.

Method: Perform comprehensive profiling of urinary metabolites by GC-TOF-MS, RP-LC-ESI-MS and HILIC-LC-ESI-MS methods, analyzing urine samples from healthy volunteers and cancer patients. Apply multivariate statistics for data mining.




USDA Agricultural Research Service
John W. Newman

USDA, ARS Western Human Nutrition Research Center (University of California, Davis)


Phase I Aim: Find potential differences between groups of individuals following high and low glycemic diet. Metabolomics pilot study. Proof of the concept.

Method: Perform blood plasma metabolite profiling by UPLC-RP-ESI-ITMS. Apply multivariate statistics for data mining.



Acknowledgments: Wei Zou, Kindra Brooks (UCD Metabolomics Core)
<http://metabolomics-core.ucdavis.edu/>

Small Molecule Biomarkers Workflow




Yale University
Yale School of Public Health
Kathleen M. McCarty

Division of Environmental Health Sciences

Phase I Aim: Find potential differences between groups of individuals with PAH exposure and without it. Among these subjects PAH-DNA adducts detected and not detected. Four groups of subjects are defined for data mining.

Method: Perform blood plasma metabolite profiling by GC-TOF-MS and UPLC-ITMS. Apply multivariate statistics for data mining.



В современном мире
наблюдается
переизбыток информации
и недостаток времени.

- Поэтому в последнее время на международных научных конференциях делают все больше стендовых докладов.



У стендового доклада есть и другие преимущества:

- 1. Представляя стендовый доклад, вы можете более свободно излагать информацию, не заботясь о времени.
- 2. Можно вступить в более тесную коммуникацию с людьми, которых интересует Ваше исследование.
- 3. Можно избежать устного доклада, если Вы страдаете дисфункцией речи

- ④ 4. Вы можете использовать этот же постер для других конференций.
- ④ 5. Стендовый доклад можно повесить в своем учебном заведении и познакомить со своими исследованием коллег, которые не смогли приехать на конференцию.
- ④ 6. Наконец, Вы можете поместить PDF своего доклада в банк постерных докладов, например на www.eposters.net, и тогда больше людей смогут прислать Вам свои комментарии.

Изготовление постера – творческий процесс, но тем не менее следует придерживаться некоторых основных принципов.

- ◎ **Читаемость**
- ◎ **Наглядность**
- ◎ **Понятность**
- ◎ Тем не менее, его прочтет большее количество людей, если Вам удастся придать ему привлекательный внешний вид!

Размер постера

- 120 см на 80 см – формат A0 (расположение листа – по вертикали или горизонта-ли).
- Постер может состоять как из одного листа формата A0, так и из нескольких листов формата A4 или A3 (в сумме не превышающих лист формата A0).

Структура

- В верхней части листа содержится следующая информация:
- * название постера,
- * ФИО автора (ов),
- * краткие сведения об авторах (уч. степень, должность и место работы),
- * контактные данные (желательно указать адрес электронной почты).



Постер (стендовый доклад) представляет собой краткое сообщение о научной работе и обычно содержит все те же разделы, что и научная статья:

- ⊙ * введение (актуальность),
- ⊙ * цель исследования,
- ⊙ * методы исследования,
- ⊙ * результаты и их обсуждение,
- ⊙ * выводы,
- ⊙ * краткий список литературы.



Иллюстрации

- * графические материалы облегчают процесс изложения доклада,
- * иллюстрации должны читаться участниками на расстоянии одного метра и далее,
- * диаграммы, рисунки и схемы предпочтительно оформлять простыми и четкими линиями,
- * заголовок постера рекомендуется печатать не менее, чем 60-м кеглем, а сам текст постера - не менее, чем 20-м.

Фотографии и рисунки выглядят выигрышнее, если обвести их узким черным контуром:



Как выбрать цвета для диаграмм и графиков?



Не используйте однотипные цвета


Используйте цвета, которые хорошо отличаются друг от друга!

Как выбрать дизайн постера?


- Используйте оформление, которое соответствует теме Вашего доклада
- Не используйте только черные и белые цвета.
- Лучше используйте пастельные цвета

Как разместить информацию на стенде?

- В европейских языках читают слева направо и сверху вниз, поэтому размещайте информацию так, чтобы читая, человек двигался от верхнего левого края к нижнему правому.




Несущественные
разделы, которые не
нужны для понимания
Вашего исследования,
такие как
благодарности, список
литературы и т.д. (хотя
некоторые люди их
читают), можно вынести
в отдельное поле внизу
доклада:




- 

Рогатые жабы: как поймать и приручить?


- При просмотрении страницы внимание в основном направлено на верхний левый угол и центр, затем правый нижний угол, затем левый нижний и правый верхний.



- Поэтому самую важную информацию лучше помещать в левый верхний угол и в центр.




- Помните, что оптический центр страницы примерно на $1/8$ выше его геометрического центра.



- Правый верхний угол лучше занять фотографиям и авторов, необходимым и эмблемами или элементом дизайна:



По способу размещения информации можно выделить два типа постеров:




С полосным расположением




и с модульным расположением.



Пример постера с полосным расположением частей:



Пример постера с модульным расположением частей:



Помните так же, что заголовки не должны совпадать по высоте:

Рогатые жабы: как поймать и приручить?

Петр Васечкин



Рогатые жабы: как поймать и приручить?

Петр Васечкин



Правильно

Неправильно

- Ширину полосы лучше сделать около 40 знаков –такой текст читается быстрее всего.
- В большинстве случаев для выделения лучше использовать *курсив*, а не *подчеркивание*

Советы и рекомендации

- * подготовьте постер и сообщение на 3-4 минуты заранее
- * при желании Вы сможете принести ноутбук и показывать с него дополнительные демонстрационные материалы (но, к сожалению, розетки в непосредственной близости от постера не будет)
- * рядом с постером можно разместить конверт с любым дополнительным материалом (визитки, лист обратной связи, копии постера и т.п.)

Помните!

- На чтение Вашего доклада не должно уходить более, чем 7 минут!
- Поэтому мы рекомендуем Вам делать текстовые отрывки не более 150 слов в длину.

Ваш текст не должен занимать более одной четверти площади слайда!



27-hydroxycholesterol is a Novel Endogenous Regulator of Estrogen Receptor Activity

Carolyn D. DuSell and Donald P. McDonnell.

Department of Pharmacology and Cancer Biology, Duke University Medical Center, Durham, NC 27710



Introduction

There are two isoforms of the estrogen receptor (ER), ER α and ER β , which upon ligand binding regulate the transcription of target genes. In general, it is thought that the proliferative effects of estrogens are mediated primarily through ER α , whereas the anti-proliferative ones are through ER β . There are three endogenous estrogen receptor agonists that activate both ER α and ER β , the most potent being 17 β -estradiol (E2). The majority of breast cancers express ER α and rely on E2 as a growth stimulus, prompting the development of therapies aimed at reducing E2 levels, such as aromatase inhibitors. Unfortunately, many breast cancers become resistant to aromatase inhibitors, but continue to rely on ER α for growth. Tumor-infiltrating macrophages (TAMs), which are associated with increased tumor growth and decreased patient survival, possess high cytochrome P450 27A1 (CYP27A1) enzymatic activity and are therefore capable of producing 27-hydroxycholesterol (27HC), which we have shown to be a novel endogenous partial agonist for ER. It is therefore possible that macrophage infiltration of a breast tumor provides an alternate estrogenic ligand to promote tumorigenic behavior. Furthermore, the production of 27HC by either TAMs and tumor epithelial cells, and the estrogenic activities of 27HC presents a potential mechanism for the development of resistance to aromatase inhibitors. Therefore, understanding the intricacies of the regulation of ER by 27HC and how this impinges on estrogen signaling will have a profound impact on estrogen-regulated biology, especially in breast cancer.

Hypothesis & Objectives

Hypothesis:


We hypothesize that 27HC is a partial agonist for ER α and ER β that may contribute to therapeutic resistance in breast cancer.

Objectives:

- ◆ Determine whether 27HC is an ER agonist, antagonist, or Selective Estrogen Receptor Modulator (SERM).
- ◆ Define the role of 27HC in regulating ER α activity in ER α -positive breast cancer cell lines.

Results


Figure 1. 27HC activates exogenous ER α and ER β



ER-negative Hela cells were transfected with pCDNA3.1myc-ER α or ER β , a 3xERE-TATA-luc reporter, and a CMV- β -gal transfection control for 24 hours. Cells were treated with vehicle or increasing concentrations of E2 or 27HC for 22-26 hours, then harvested and assayed for luciferase and β -gal expression. Data is presented as mean \pm SEM for three independent triplicate experiments.


Results

Figure 2. 27HC regulates endogenous ER α target genes




ER α -positive MCF-7 cells were plated in phenol-free media containing 8% charcoal-stripped FBS at 2×10^5 cells/ml. After 48 hours, the cells were treated with vehicle or ligand. At the indicated time, cells were harvested for RNA isolation and ER target gene expression was analyzed by RT-PCR. Threshold cycle values were normalized to the housekeeping gene 36B4. Data was then normalized to vehicle and is presented as mean \pm SEM for three independent experiments.

Figure 3. 27HC treatment leads to recruitment of ER α at the estrogen response element in the pS2 promoter



MCF-7 cells were plated in phenol-free media containing 8% charcoal-stripped FBS at 3.5×10^5 cells/ml. After 72 hours, the cells were treated with vehicle, 100nM E2, or 10 μ M 27HC for 45 minutes. After cross-linking DNA and protein, protein and associated chromatin was harvested and subjected to immunoprecipitation with either IgG or ER α antibody. Following immunoprecipitation, cross-linking was reversed and DNA harvested for analysis by RT-PCR. Data is presented as the percent of DNA immunoprecipitated with ER α compared to the total amount of DNA present in the input.

Figure 4. 27HC stimulates proliferation in ER α -positive breast cancer cells




MCF-7 cells were plated in phenol-free media containing 8% charcoal-stripped FBS at 5×10^4 cells/ml. After 48 hours, the cells were seeded and the media was replaced with serum-free media for 24 hours. Cells were then treated with vehicle, E2, or 27HC for 48 hours. Treatment was replenished after 48 hours. On day 5, the increase in cell number was assessed using a DNA dye. Data is presented as the mean \pm SEM for three independent triplicate experiments.

Carolyn D. DuSell
Ph.D. Student, Department of Pharmacology and Cancer Biology
Duke University Medical Center
Durham, NC 27710
Phone: 919.286.2111
Email: cdsell@duke.edu


Results

Figure 5. 27HC induces a unique conformation of ER α and ER β



ER-negative HepG2 cells were transfected with ER-interacting peptides fused to Gal4 to use as conformational probes of the receptor structure. VP16-ER α or ER β , a 5xGAL4-TATA-luc reporter, and a CMV- β -gal transfection control were co-transfected with the peptides for 24 hours. Cells were subsequently treated with vehicle, 10 μ M 27HC, 100nM E2, 100nM 4OHT, or 100nM ICI for 22-26 hours. Cells were then harvested and assayed for luciferase and β -gal expression. Data is presented as the mean of one representative triplicate experiment.

Figure 6. 27HC allows for differential peptide recruitment to ER α and ER β



HepG2 cells were transfected with VP16-ER α or ER β , Gal4 peptide, 5xGAL4-TATA-luc reporter, and CMV- β -gal transfection control for 24 hours. Cells were subsequently treated with vehicle or increasing concentrations of 27HC or E2 for 22-26 hours. Cells were then harvested and assayed for luciferase and β -gal expression. Data is represented as the mean of one representative triplicate experiment.

Conclusions

- ◆ 27HC is a novel partial agonist that regulates both ER α and ER β .
- ◆ 27HC stimulates proliferation in an ER α -positive breast cancer cell line.
- ◆ 27HC induces a unique conformation of both ER α and ER β that allows for differential recruitment of ER-interacting peptides.

Future Directions

- ◆ Determine how 27HC influences breast tumor growth.
- ◆ Establish which cells produce 27HC within the tumor microenvironment and determine whether this ligand acts in an autocrine or paracrine manner.

Acknowledgements

This project is funded in part by the Department of Defense Predoctoral Traineeship Award BC050609.

Как распечатать стендовый доклад?

- Мы рекомендуем Вам создать макет Вашего постера в одной из графических программ. Программа PosterGenius создана специально для изготовления научных постеров. <http://www.postergenius.com>
- Затем можно разделить постер в PDF
- Однако большинство исследователей пользуется такими программами, как:
- InDesign, QuarkXPress, Pagemaker, Illustrator, CorelDRAW, Freehand, Powerpoint и др.
- А затем отправь постер нам и мы распечатаем его)))

- Рекомендации по оформлению постера (на английском языке) можно прочитать тут -
<http://colinpurrington.com/tips/academic/posterdesign> (ссылка заимствована на сайте
<http://virtualcoglab.ru/poster.html>)
- В стандартных приложениях Windows существует программа Microsoft Office Publi-sheer, позволяющий самостоятельно делать постеры. Существуют и другие про-граммы.